



CanadaPetCare.com

Owned By Veterinarian

BRAVECTO CHEWABLES FOR DOGS



Recommended for dogs only.

Used for the long-lasting protection from fleas and ticks on dogs and puppies 6 months of age.

INDICATIONS

- Indicated for the treatment and prevention of flea (*Ctenocephalides felis*) infestations for three months.
- Treatment and control of paralysis tick (*Ixodes holocyclus*) for four months.

- Treatment and control of brown dog tick (*Rhipicephalus sanguineus*) for eight weeks.
 - One chew provides consistent and long lasting efficacy against both fleas and ticks.
 - Treats and controls tick infestations including black-legged tick, American dog tick.
 - Indicated for the treatment and control of lone star tick infestations for 8 weeks.
 - Controls flea and tick infestations in dogs and puppies 6 months of age and older, and weighing 4.4 pounds or greater.
-

COMPOSITION

Fluralaner

RESIDUAL EFFECT

Fluralaner in Bravecto takes effect quickly and starts killing fleas – both adult and juvenile within 2 hours, and ticks within 12 hours.

WARNING

- For veterinary use only.
- Keep this drug out of reach of children and other animals.

- Keep the product in the original package until in use.
 - Do not eat, drink or smoke while handling the product.
 - Wash hands thoroughly with soap and water immediately after use.
-

PRECAUTIONS

- Do not administer in puppies less than 6 months of age.
 - Bravecto is not effective against ticks beyond 8 weeks after dosing.
-

DOSAGE

- Bravecto should be administered orally with a minimum dose is 11.4 mg/lb. body weight as standard.

Weight of dog (lbs)	Tablets to use per dose
4.4 – 9.9	One yellow box tablet (112.5 mg)
10 – 22	One orange box tablet (250 mg)
22.1 – 44	One green box tablet (500 mg)
44.1 – 88	One blue box tablet (1000 mg)
88.1 – 123	One pink box tablet (1400 mg)
Over 123	Use an appropriate combination.

METHOD OF APPLICATION

- Administer Bravecto orally as a single dose every 12 weeks according to dosage schedule.
 - Give each dose of Bravecto with food to reduce the risk of vomiting.
 - Can be given any time of the year and can continue year round without interruption.
-

STORAGE AND DISPOSAL

- Do not store above 86°F (30°C).
 - Keep away from direct sunlight.
 - Avoid keeping near food products.
-

PESTICIDE DISPOSAL

Dispose of empty package by wrapping with paper and putting in garbage.

SIDE EFFECTS

In rare cases, a dog treated with Bravecto may show signs of vomiting.

CONTRAINDICATIONS

There are no known contraindications for the use of the product.

FIRST AID FOR HUMANS

- If product is swallowed or gets in mouth, do not induce vomiting. Contact a poison information center or call a doctor.
- In case of skin contact, wash with lukewarm water for 5 minutes.
- In unlikely event of eye contact, immediately wash eyes with water thoroughly.

Manufactured by

Merck Animal Health



Visit our website at -

<https://www.canadapetcare.com/bravecto-for-dogs/flea-and-tick-control-treatment-278.aspx>

DISCLAIMER

The information provided in this file is for informational purpose only. Please consult your veterinarian before administering the product to your pet.

© 2017 Canadapetcare.com - All Rights Reserved.