



**CanadaPetCare.com**  
Owned By Veterinarian

---

## **BRAVECTO SPOT ON FOR CATS**



For use on cats only.

For use on cats and kittens 6 months of age and older and weighing 2.6 lbs. or greater.

---

## **INDICATIONS**

- Indicated for the treatment and prevention of flea infestations.
- Treatment and control of black-legged tick infestations for 12 weeks.
- Helps prevent American dog tick infestations.
- Aids to reduce signs of flea allergy dermatitis.

## **COMPOSITION**

Fluralaner

---

## **RESIDUAL EFFECT**

Bravecto starts killing fleas about 8 hours and ticks about 12 hours of administration. It remains effective for about 12 weeks.

---

## **WARNING**

- Not for human use. Keep away all drugs out of reach of children.
  - Do not allow children to touch the application site until dry.
  - Keep the product in the original package until in use.
  - Do not eat, drink or smoke while handling the product.
  - Avoid contact with skin and eyes.
  - Wash hands and contacted skin thoroughly with soap and water immediately after use of the product.
  - Keep the product away from direct open flame, heat and sparks.
-

## **PRECAUTIONS**

- Avoid oral ingestion. For topical use only.
  - The safety of Bravecto has not been established in breeding, pregnant and lactating cats.
  - Do not use in cats with a history of neurologic abnormalities.
- 

## **DOSAGE**


Bravecto should be administered topically as a single dose every 12 weeks with a minimum dose of 18.2 mg/lbs. (40 mg/kg) body weight.

---

## **METHOD OF APPLICATION**

- Open the pack prior to treatment and remove the tube.
- Hold the tube at the upright position. Rotate the cap in a clockwise direction or anti-clockwise direction in one full turn.
- The cap is designed to stay on the tube for dosing and should not be removed.
- When a “click” sound is heard, the tube is ready to open and apply.
- This ensures seal is broken and the solution is ready to apply.
- Make the cat standing or lying with its back horizontal during the application.
- Part the fur until skin is visible at the administration site.

- Place the tube tip vertically horizontal against the skin at the base of the skull of the cat.



- Squeeze the tube and gently apply the entire contents of Bravecto directly to the skin at the base of the skull of the cat.
  - Avoid applying an excessive amount of the solution that could cause solution to drip off the cat.
  - If to avoid run off, second spot is needed, then apply the second spot slightly behind the first spot.
  - Treat the cat at an interval of 12 weeks.
  - Treatment with Bravecto can begin any time of the year and can continue year round with without interruption.
- 

## **STORAGE AND DISPOSAL**

- Do not store above 86°F (30°C).
  - Store in the original package in order to protect from moisture.
  - The pouch should only be opened immediately prior for use.
- 

## **PESTICIDE DISPOSAL**

Dispose of empty containers by wrapping with paper and putting in garbage.

## **SIDE EFFECTS**

In rare cases, a mild short-lived skin reaction at the application site occurs.

---

## **CONTRAINDICATIONS**

There are no known contraindications for the use of the product.

---

## **FIRST AID**

In case of eye contact, rinse off immediately with water.

---

Manufactured by  
Merck Animal Health



**Visit our website at –**

<https://www.canadapetcare.com/bravecto-spot-on-for-cats/flea-and-tick-control-treatment-294.aspx>

## **DISCLAIMER**

The information provided in this file is for informational purpose only. Please consult your veterinarian before administering the product to your pet.

© 2017 Canadapetcare.com - All Rights Reserved.