

FRONTLINE TOP SPOT FOR DOGS



For use on dogs only. Do not use on cats or rabbits.

For use on puppies and dogs from 8 weeks of age.

INDICATION

- A topical spot-on treatment that treats and controls flea infestation, brown dog tick, American dog tick, lone star tick, deer ticks and chewing lice on dogs and puppies.
- Assists in treating Flea Allergy Dermatitis (FAD).
- Completely waterproof treatment.
- Contains Fipronil as an active ingredient which inhibits the regulatory channels resulting in continuous excitement of nervous system of the dog thus causing death of adult fleas and ticks.
- Controls Sarcoptic mange infestations.
- Fast acting and long lasting.
- Kills adult 100% fleas within 24 hours of administration.

- Kills newly emerged adult fleas before they lay eggs.
 - Recommended for breeding, pregnant or lactating dogs.
 - Prevents paralysis ticks for two weeks and brown ticks for full month.
 - Very easy to apply.
-

COMPOSITION

Fipronil	9.70%
----------	-------

RESIDUAL EFFECT

A monthly spot-on treatment that kills adult fleas along with biting lice on dogs. It kills 100% adult fleas within 24 hours of administration along with prevention of paralysis ticks for two weeks and brown ticks for the full month. It prevents sarcoptic mange and aids in treating Flea Allergy Dermatitis (FAD).

WARNING

- Never administer this treatment on dogs that are allergic to the ingredient present in it.
 - Ensure that it is not ingested by the dog. Use it only externally.
 - Never use this treatment on puppies younger than 8 weeks of age.
 - Do not use this treatment in conjunction with other pesticides on dogs.
 - Keep the product away from the reach of children and other pets and do not allow them to get in contact with the treated dog till the application site dries out completely.
-

PRECAUTIONS

- The safety of this product on debilitated or aged dog has not been verified yet. So, consult with your veterinarian before administering this treatment



on him/her.


DOSAGE

DOG WEIGHT	DOSAGE
up to 22 lbs.	1 pipette of 0.67 ml
23 lbs. to 44 lbs.	1 pipette of 1.34 ml
45 lbs. to 88 lbs.	1 pipette of 2.68 ml
over 89 lbs. to 132 lbs.	1 pipette of 4.02 ml
over 132 lbs.	2 pipettes of 2.68 ml

METHOD OF APPLICATION

- Use as directed by your veterinarian.
- Recommended for puppies and dogs at least 8 weeks of age.

Step-1		Hold the pipette in an erect position. Tap the thin part of the pipette and check that the contents are inside the principal body of the pipette. Unlock the snap-off top of the product pipette along the graded line.
Step-2		Divide the coat between the shoulder blades till the skin can be seen.

Step-3		Keep the tip of the pipette on the dog's skin and compress softly at a single or multiple spots to empty its content on the skin of the dog.
---------------	---	--

- This treatment needs to be repeated on a monthly basis on the same date.
-

STORAGE AND DISPOSAL

- **Storage:** Store the product at a room temperature below 30 degrees Celsius. Store unused product in original container only, out of reach of children and pets.
 - **Disposal:** Mask blank container in paper and place in garbage
-

PESTICIDE DISPOSAL

- Do not contaminate water, food, or feed by storage and disposal.
 - Never empty product in indoor or outdoor drain.
-

CONTAINER DISPOSAL

- **If the container is empty:** Never re-use or refill the container. Keep it in trash or offer for recycling if the option is available.
 - **If the container is wholly filled:** Contact your local solid waste agency for disposal instructions. Never keep the new product near any indoor or outdoor drain.
-

FIRST AID

Keep the product container or label with you while calling a poison control center or doctor or while going for the treatment:

- **If the treatment is swallowed:** Contact poison contact center or doctor

quickly for getting their advice. If the person is able to swallow, give him/her a sip of glass of water. Never induce vomiting until and unless it has been recommended by the poison center or doctor. If the person is unconscious do not give anything by mouth.

- **If the treatment gets into the eyes:** Keep the eye open and rinse slowly and softly with water for 15 to 20 minutes. If the person is wearing contact lenses, remove it after 5 minutes and continue rinsing the eye. Call the doctor or poison control center for advice.
- **If the treatment affects the skin:** Wash the spot with plenty of soap and water. Take the help of a doctor in case if the affected being is a person. For the dog contact your veterinarian immediately.

SIDE EFFECTS

In certain dogs, there may be temporary redness or itchiness at application site. If the application site is licked by the dog he might drool before the treatment dries. If it persists for several hours, bathe the dog and contact your veterinarian.

Manufactured by
Merial



Visit our website at

<https://www.canadapetcare.com/frontline-top-spot-for-dogs/flea-and-tick-control-treatment-11.aspx>

DISCLAIMER

The information provided in this file is for informational purpose only. Please consult your veterinarian before administering the product to your pet.

© 2016 CanadaPetCare.com - All Rights Reserved.

